



# Orthopaedic & Sports Medicine

**Jamie L. Friedman, M.D.**

---

## **Achilles Tendon Repair Rehab Protocol**

### **POST OPERATIVE MANAGEMENT**

- Posterior Splint (short leg, neutral ankle) / Non-Weight-Bearing with crutches

### **Week 2 – (After first post-op visit)**

- Discontinue splint use and transition to CAM walking boot with heel lift to maintain plantar flexion position
- Advance weight bearing to full / wean from crutches over 1-2 weeks
- Cryotherapy (Jobst compression/Cryocuff/Ice pack)
- Avoid passive heel-cord stretching
- Active dorsiflexion only
- Size of heel wedge may be decreased by therapist as passive foot position becomes comfortable at 90deg

### **Week 2-6:**

- Progressive weightbearing ambulation with walking boot to FWB at 3-4 weeks
- BAPs "Level 1-2-3" (seated PWB)
- Active range-of-motion dorsiflexion/plantar flexion unloaded
- Heel-Toe ambulation in boot
- Stationary cycling with minimal resistance (anterior foot placement)
- Flat-Footed standing balance/proprioception activities

### **Week 6-12:**

- Discontinue walking boot, use heel cups for 2 weeks as needed
- Progressive resistance Theraband ankle strengthening exercises
- Intrinsic foot muscle strengthening exercises
- Single Leg standing balance activities
- Active bilateral heel raises
- Initiate passive heel cord stretching
- Bilateral to unilateral standing, heel raise exercises
- Stationary cycling with progressive resistance (standard foot placement)
- Progressive depth (30° to 90° knee flexion) flat-footed mini-squats
- Progressive retrograde treadmill ambulation
- "Flat-Footed" single leg balance
- BAPs "Level 1-2-3" (seated PWB progressed to FWB standing)
- "On-Toes" standing balance/proprioception activities on mini-trampoline
- Progressive duration/velocity Stairmaster ambulation
- Lateral sliding board maneuvers

### **3-6 Months:**

- Stationary jogging and jumping on mini-trampoline
- Straight ahead jog-to-run progression on a level surface
- Progressive acuity cutting agility maneuvers / Large-to-small Figure of 8 agility maneuvers
- Sport/Position specific functional activities
- Running and agility maneuvers on progressive incline/decline surfaces
- Discharged to normal activities when cleared by doctor

### **Criteria for discharge:**

1. Full, pain free range of motion
2. Strength is equal bilaterally
3. Has met specific functional/activity goals
4. Has been cleared by physician